

16

□□□□□□□□□□□□□□**TSC**□

□ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □

16

16.....ADHD

Image



- 46% of children with ADHD have comorbid anxiety disorders, 25% have comorbid depression, 40% have comorbid conduct disorder, 25% have comorbid oppositional defiant disorder, 30% have comorbid tic disorders, and 60% have comorbid specific phobias.
- 46% of children with ADHD have comorbid anxiety disorders, 25% have comorbid depression, 40% have comorbid conduct disorder, 25% have comorbid oppositional defiant disorder, 30% have comorbid tic disorders, and 60% have comorbid specific phobias.

□ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □



01

01

□□□□



□□□□1□6□□

□□□□1□6□□

□□□□□

□□□□



□□□□7□12□□

□□□□7□12□□

□□□□□

□□□□



□□□□13□19□□

□□□□13□19□□

□□□□□

□□□□



□□□□**20**□□

□□□□**20**□□

□□□□

□□□□



□□□□

□□□□

□□□□

□□□□

□□□□□□□□□□□□□□

□□□□□□

Source URL: *https://www.pro.novartis.com/jp-ja/products/afinitor/tsc/age_03*