

5,007 16.01 /1,000 295
/18,426 2,829 19.07 /1,000 190 /9,963 0.84 95
0.62 1.13 IgE

2019 12 1

1

150mg 2.8% 2/71
5.4% 4/74 300mg^{2,3}
284 53 18.7% 26 9.2%
24 8.5% 20 7.0% 14 4.9%
13 4.6% 12 4.2%
38 3 7.9% 2 5.3%

2

Churg-Strauss^{2,3}
¹

3

$\leq 100 \times 10^9/L$ ^{2,3}
1304 1307 0.4 2/284^{4,5}

4

300mg 1.4 1/73
1.4 1/74^{2,3}
8 3,342 2.69
/1,000 5 /1,856 2,895 2.38 /1,000 4 /1,680 1.13 95
0.24 5.71 2 1 3
1 5 5,007 7.52 /1,000
115 /15,286 2,829 5.12 /1,000 51 /9,963

1.32 95 0.91 1.91¹⁰

¹⁰

※

1 2019 12 1

2 20170085, ,

3 Hide, M. et al.: J. Dermatol. Sci. 87(1), 70, 2017 20170808, /
4

4 Ohta, K. et al.: Respirology 14(8), 1156, 2009 XOLJ00021, 2

5 Ohta, K. et al.: Allergol. Int. 59(2), 167, 2010 XOLM00054, 3

2306

	144	
	13	
	9.0	
	300mg N73	150mg N71
	0 0.0	0 0.0
	0 0.0	0 0.0
	0 0.0	0 0.0
	0 0.0	0 0.0
	1 1.4	0 0.0
	1 1.4	0 0.0
	0 0.0	1 1.4

□□□□□□	0□0.0□	1□1.4□
□□□□□□	0□0.0□	1□1.4□
□□□□□□□□	0□0.0□	0□0.0□
□□□□□□	0□0.0□	1□1.4□
□□□□□□	0□0.0□	0□0.0□
□□□□□□□□	0□0.0□	0□0.0□
□□□□□□□□□□	1□1.4□	1□1.4□
□□□□	1□1.4□	1□1.4□
□□□□	0□0.0□	0□0.0□
□□□□□□	0□0.0□	0□0.0□
□□□□□□	3□4.1□	1□1.4□
□□	2□2.7□	1□1.4□
□□□	1□1.4□	0□0.0□
□□□□□□□□	1□1.4□	0□0.0□
□□	1□1.4□	0□0.0□
□□□□□□□□□□□□□□	2□2.7□	1□1.4□
□□□□□□	1□1.4□	0□0.0□
□□□□□	1□1.4□	0□0.0□
□□□□□	0□0.0□	1□1.4□
□□□□□□□□□□□□	1□1.4□	3□4.2□
□□	1□1.4□	0□0.0□
□□□□□□□□□□	0□0.0□	1□1.4□
□□□□□	0□0.0□	1□1.4□
□□□□□	0□0.0□	0□0.0□
□□□	0□0.0□	1□1.4□

□□□□□□□□

□□□□□□□□

□□□□□□

□□□□□□□□□□□□

□□□□□□□□

□□□□□

□□□□

□□□□□□

□□□□

PFS□□□□

□□□□□□□

□□□□□□□□□□□□

□□□□□□

□□□□□□□□□□□□□□□□□□

Image



□□□□□□□□□□□□

Source URL: https://www.pro.novartis.com/jp-ja/products/xolair/urticarial/point_04