

Novartis Pharmaceuticals Corporation

Novartis Pharmaceuticals Corporation, 235 East 41st Street, New York, NY 10017

Novartis

Clozapine (Clozaril) Tablets

Section	Text
Warnings	See important information about Clozapine (Clozaril) Tablets on the adjacent page.
Contraindications	Clozapine (Clozaril) Tablets are contraindicated in patients with a history of myocardial infarction, stroke, or other conditions that may be exacerbated by the effects of Clozapine (Clozaril) Tablets. Clozapine (Clozaril) Tablets are also contraindicated in patients with a history of bone marrow depression or other conditions that may be exacerbated by the effects of Clozapine (Clozaril) Tablets.
Precautions	Clozapine (Clozaril) Tablets may cause drowsiness, dizziness, and blurred vision. Patients should avoid driving or operating machinery until they know how Clozapine (Clozaril) Tablets affect them. Clozapine (Clozaril) Tablets may also cause changes in heart rate and blood pressure. Patients should be monitored for these effects. Clozapine (Clozaril) Tablets may also cause changes in white blood cell counts. Patients should be monitored for these effects.
Adverse Reactions	Common adverse reactions include drowsiness, dizziness, blurred vision, and changes in heart rate and blood pressure. Other adverse reactions include changes in white blood cell counts, bone marrow depression, and other conditions that may be exacerbated by the effects of Clozapine (Clozaril) Tablets.
Drug Interactions	Clozapine (Clozaril) Tablets may interact with other medications, including antidepressants, antipsychotics, and other drugs that may affect the heart or blood pressure. Patients should inform their healthcare provider of all medications they are taking.
Use in Specific Populations	Clozapine (Clozaril) Tablets should be used with caution in patients with liver or kidney disease. Clozapine (Clozaril) Tablets should also be used with caution in pregnant and breastfeeding women.
Overdosage	Overdosage of Clozapine (Clozaril) Tablets may cause drowsiness, dizziness, blurred vision, and changes in heart rate and blood pressure. Patients should seek medical attention if they experience these symptoms.
How to Take Clozapine (Clozaril) Tablets	Clozapine (Clozaril) Tablets should be taken as directed by your healthcare provider. Do not take more than the prescribed amount. Do not stop taking Clozapine (Clozaril) Tablets without consulting your healthcare provider.

Novartis Pharmaceuticals Corporation

Novartis Pharmaceuticals Corporation, 235 East 41st Street, New York, NY 10017

Novartis

<mailto:clozaril.remote@novartis.com>

Novartis 0800-222-8814

Novartis 900-1730

Novartis

Novartis 0120-003-293

Novartis 900-1730

Novartis

Source URL: <https://www.pro.novartis.com/jp-ja/products/clozaril/clinical/13>