

NEN

NEN

NEN NEN NEN

NEN NEN NEN

NEN

NEN VIP 1 NEN 1

NEN

NEN NEN NEN

1 NEN

^{1,2}	<ul style="list-style-type: none">92% 51%38%43% 23%	NIPHS

<p> Zollinger-Ellison syndrome^{1,2} </p>	<ul style="list-style-type: none"> • 26-98% • 0-56% • 71-93% • 17-73% • 8-75% 	<p> Zollinger-Ellison syndrome </p>
<p> VIPoma^{1,2} </p>	<ul style="list-style-type: none"> • 22-90% • 14-18% • 54-90% 	<p> VIPoma </p>
<p> VIPoma WDHA^{1,2} </p>	<ul style="list-style-type: none"> • 89-100% • 67-100% • 44-100% • 14-33% 	<p> VIPoma WDHA </p>
<p> NIPHS^{1,3} </p>	<ul style="list-style-type: none"> • 58-100% • 45-96% • 3-18% • 14-41% 	<p> NIPHS </p>

NIPHS: non-insulinoma pancreatogenous hypoglycemia syndrome, WDHA: watery diarrhea hypokalemia achlorhydria

1 Ito T, Jensen R. Expert Opin Pharmacother. 2021; 22: 685-693.

2 . 2014; 111: 2263-2271.

3 Khan MS, et al. World J Gastroenterol. 2020; 26: 4537-4556.

NEN diagnosis: US, CEUS, CT, MRI, SRS, SASI test, PET/CT, FDG-PET/CT.

NEN

	<ul style="list-style-type: none"> • CT/MRI • US • EUS/EUS-FNA • SASI test

